

§ 172.800

102™,” dated October 9, 2003, and printed by Intralytix, Inc., is incorporated by reference. The Director of the Office of the Federal Register approves this incorporation by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. You may obtain a copy from the Office of Food Additive Safety (HFS-200), Center for Food Safety and Applied Nutrition, Food and Drug Administration, 5100 Paint Branch Pkwy., College Park, MD 20740, or you may examine a copy at the Center for Food Safety and Applied Nutrition’s Library, 5100 Paint Branch Pkwy., College Park, MD 20740, or at the National Archives and Records Administration (NARA). For information on the availability of this material at NARA, call 202-741-6030, or go to: http://www.archives.gov/federal_register/code_of_federal_regulations/ibr_locations.html.

(2) The mean phage titer of each monophage in the additive is 1×10^9 plaque forming units (PFU)/ml. The analytical method for determining phage titer entitled “Method to Determine Lytic Activity/Phage Titer,” dated November 6, 2001, and printed by Intralytix, Inc., is incorporated by reference. Copies are available at locations cited in paragraph (b)(1) of this section.

(3) The phages present in the preparation must not contain a functional portion of any of the toxin-encoding sequences described in 40 CFR 725.421(d). No sequences derived from genes encoding bacterial 16S ribosomal RNA are present in the complete genomic sequence of the phages.

(4) *L. monocytogenes* toxin, listeriolysin O (LLO), is not greater than 5 hemolytic units (HU)/ml. The analytical method for determining LLO entitled “Quantitation of Listeriolysin O Levels in LMP-102™,” dated September 27, 2004, and printed by Intralytix, Inc., is incorporated by reference. Copies are available at locations cited in paragraph (b)(1) of this section.

(5) The additive is negative for *L. monocytogenes*. The modified version of the U.S. Department of Agriculture’s method for determining *L. monocytogenes* entitled “LMP-102™ *Listeria monocytogenes* Sterility Test-

21 CFR Ch. I (4–1–14 Edition)

ing,” dated May 24, 2004, and printed by Intralytix, Inc., is incorporated by reference. Copies are available at locations cited in paragraph (b)(1) of this section.

(6) The additive is negative for gram-positive and gram-negative bacteria capable of growing in commonly used microbiological media (e.g., Luria-Bertani (LB) medium), including *Escherichia coli*, *Salmonella* species and coagulase-positive *Staphylococci*, as determined by the “Method to Determine Microbial Contamination,” dated July 11, 2003, and printed by Intralytix, Inc., is incorporated by reference. Copies are available at locations cited in paragraph (b)(1) of this section.

(7) Total organic carbon (TOC) is less than or equal to 36 mg/kg. The analytical method for determining TOC entitled “Determination of Total Organic Carbon by Automated Analyzer,” dated March 30, 2001, and printed by Intralytix, Inc., is incorporated by reference. Copies are available at locations cited in paragraph (b)(1) of this section.

(c) *Conditions of use.* The additive is used in accordance with current good manufacturing practice to control *L. monocytogenes* by direct application to meat and poultry products that comply with the ready-to-eat definition in 9 CFR 430.1. Current good manufacturing practice is consistent with direct spray application of the additive at a rate of approximately 1 mL of the additive per 500 cm² product surface area.

[71 FR 47731, Aug. 18, 2006]

Subpart I—Multipurpose Additives

§ 172.800 Acesulfame potassium.

Acesulfame potassium (CAS Reg. No. 55589–62–3), also known as acesulfame K, may be safely used as a general-purpose sweetener and flavor enhancer in foods generally, except in meat and poultry, in accordance with current good manufacturing practice and in an amount not to exceed that reasonably required to accomplish the intended technical effect in foods for which standards of identity established under section 401 of the Federal Food, Drug, and Cosmetic Act do not preclude such use, under the following conditions:

(a) Acesulfame potassium is the potassium salt of 6-methyl-1,2,3-oxathiazine-4(3H)-one-2,2-dioxide.

(b) The additive meets the following specifications:

(1) Purity is not less than 99 percent on a dry basis. The purity shall be determined by a method titled "Acesulfame Potassium Assay," which is incorporated by reference. Copies are available from the Center for Food Safety and Applied Nutrition (HFS-200), Food and Drug Administration, 5100 Paint Branch Pkwy., College Park, MD 20740, or available for inspection at the National Archives and Records Administration (NARA). For information on the availability of this material at NARA, call 202-741-6030, or go to: http://www.archives.gov/federal_register/code_of_federal_regulations/ibr_locations.html.

(2) Fluoride content is not more than 30 milligrams per kilogram, as determined by method III of the Fluoride Limit Test of the Food Chemicals Codex, 7th ed. (2010), p. 1151, which is incorporated by reference. The Director of the Office of the Federal Register approves this incorporation by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. You may obtain copies from the United States Pharmacopeial Convention, 12601 Twinbrook Pkwy., Rockville, MD 20852 (Internet address <http://www.usp.org>). Copies may be examined at the Food and Drug Administration's Main Library, 10903 New Hampshire Ave., Bldg. 2, Third Floor, Silver Spring, MD 20993, 301-796-2039, or at the National Archives and Records Administration (NARA). For information on the availability of this material at NARA, call 202-741-6030 or go to: <http://www.archives.gov/federal-register/cfr/ibr-locations.html>.

(c) If the food containing the additive is represented to be for special dietary uses, it shall be labeled in compliance with part 105 of this chapter.

[53 FR 28382, July 28, 1988, as amended at 57 FR 57961, Dec. 8, 1992; 59 FR 61540, 61543, 61545, Dec. 1, 1994; 60 FR 21702, May 3, 1995; 63 FR 36362, July 6, 1998; 68 FR 75413, Dec. 31, 2003; 78 FR 71464, Nov. 29, 2013]

§ 172.802 Acetone peroxides.

The food additive acetone peroxides may be safely used in flour, and in bread and rolls where standards of identity do not preclude its use, in accordance with the following prescribed conditions:

(a) The additive is a mixture of monomeric and linear dimeric acetone peroxide, with minor proportions of higher polymers, manufactured by reaction of hydrogen peroxide and acetone.

(b) The additive may be mixed with an edible carrier to give a concentration of: (1) 3 grams to 10 grams of hydrogen peroxide equivalent per 100 grams of the additive, plus carrier, for use in flour maturing and bleaching; or (2) approximately 0.75 gram of hydrogen peroxide equivalent per 100 grams of the additive, plus carrier, for use in dough conditioning.

(c) It is used or intended for use: (1) In maturing and bleaching of flour in a quantity not more than sufficient for such effect; and (2) as a dough-conditioning agent in bread and roll production at not to exceed the quantity of hydrogen peroxide equivalent necessary for the artificial maturing effect.

(d) To insure safe use of the additive, the label of the food additive container and any intermediate premix thereof shall bear, in addition to the other information required by the act:

(1) The name of the additive, "acetone peroxides".

(2) The concentration of the additive expressed in hydrogen peroxide equivalents per 100 grams.

(3) Adequate use directions to provide a final product that complies with the limitations prescribed in paragraph (c) of this section.

§ 172.804 Aspartame.

The food additive aspartame may be safely used in food in accordance with good manufacturing practice as a sweetening agent and a flavor enhancer in foods for which standards of identity established under section 401 of the act do not preclude such use under the following conditions:

(a) Aspartame is the chemical 1-methyl *N*-L- α -aspartyl-L-phenylalanine (C₁₄H₁₈N₂O₅).